

LOCAL PLANTS OF THE INNER COASTAL PLAIN
OF SOUTHEASTERN VIRGINIA

M. L. FERNALD

(Continued from page 415)

ZORNIA BRACTEATA (Walt.) Gmel. SOUTHAMPTON COUNTY: "Plentiful in a dry sandy field at Franklin,"¹ 1893, *Heller*, no. 1029, "new to northern range." ISLE OF WIGHT COUNTY: forming broad carpets, border of sandy yellow pine and oak woods north of Walters, *F. G. & L.*, no. 6624. See p. 354.

STYLOSANTHES BIFLORA (L.) BSP., var. HISPIDISSIMA (Michx.) Pollard & Ball. ISLE OF WIGHT COUNTY: dry sandy pine and oak woods about 1 mile southeast of Zuni, *F. & L.*, no. 6241; similar habitat near Walters, *F. & L.*, no. 6242. See p. 350.

LATHYRUS VENOSUS Muhl. SUSSEX COUNTY: dry sandy hickory and oak woods, Burt, *F. & L.*, no. 6244.

This species, like several others with it, seen only once on the Coastal Plain.

*LATHYRUS HIRSUTUS L. HENRICO COUNTY: roadside, entrance to University Heights, Westhampton, *F. L. & S.*, no. 5814.

A European species becoming established in several parts of this country.

*GALACTIA MACREEI M. A. Curtis in *Bost. Journ. Nat. Hist.* i. 120 (1837). *G. pilosa*, β . *Macreei* (M. A. Curtis) Torr. & Gray, *Fl. N. Am.* i. 287 (1838), as β . "*Macraei*." PRINCESS ANNE COUNTY: climbing 1–2.5 m., thickets and woods, Dam Neck, *F. & L.*, no. 3978; wet thicket near outlet of Rainey's Pond, Sand Bridge, *F. & L.*, no. 3980; both distributed as *G. volubilis* (L.) Britton. Corolla delicate pink, with purple center.

Galactia Macreei is one of three species which have been passing as *G. volubilis*. Linnaeus based his *Hedysarum volubile* L. *Sp. Pl.* 750 (1753) upon the plate of *H. trifolium scandens* of Dillenius, *Hort. Elth.* 173, t. 143, which, in absence of other material, must stand as type. The Dillenian plate at once suggests *G. mollis* Michx. (1803), as recognized by Miss Vail in *Bull. Torr. Bot. Cl.* xxii. 507 (1895). In the latter species, however, the dense pubescence of stem, petioles, and peduncles ascends or points upward; in *G. volubilis* and its allies it is reflexed, as in the Dillenian plate. The distinctions between *G. Macreei* and *G. volubilis* follow.

G. VOLUBILIS. Pubescence of stem, etc., loosely spreading or loosely reflexed: leaflets oval to oval-oblong, the larger 1–3 cm. broad: peduncles

¹ See *Heller*, l. c. 23 (1894).

and flowering rachises stiff, pilose, in the best-developed racemes 3–11 cm long, floriferous nearly to base, the true peduncles only 1 mm.–3.5 cm. long; the groups of flowers 0.5–2 cm. apart: full-grown flower-buds, just before expansion, slightly curved; the beak (sepal-tips) about 1/3 length of body: calyx spreading-pilose, 4–5.5 mm. long: the basal bracts ovate: corolla pink, essentially unicolorous; keel-petals 6–7 mm. long (in var. *mississippiensis* Vail 6–10 mm.): legumes densely spreading-pilose, 2–5.5 cm. long.—Dry thickets and borders of woods, Florida to Texas, north to Long Island, Indiana, Missouri and Kansas.

G. MACREEI (isotype in Gray Herb.). Pubescence of stem, etc., minute, retrorsely strigillose: leaflets oblong, the larger 0.5–2 cm. broad: peduncles and flexuous rachises filiform, retrorsely strigillose or glabrous, in the best-developed racemes 0.7–3 dm. long, flowering only well above the base; the true peduncles 3–7 cm. long; the groups of flowers 1.5–4 cm. apart: full-grown flower-buds with longer and more falcate beak: calyx subappressed-pilose, 6–10 mm. long, its basal bracts linear- or lance-subulate: corolla pink, with deep purple center; the keel-petals 9–10 mm. long: legumes minutely strigose, 3–7 cm. long.—Damp or wet thickets, pond-margins and low woods, Florida to Texas, north on Coastal Plain to southeastern Virginia.

When Miss Vail, l. c., said "*Galactia Macreei*, the type specimen of which is preserved in Herb. Columbia College, is merely a very slender filiform-racemed variation" of *G. volubilis*, she evidently did not make close comparisons of the details. At that time she treated *G. pilosa*, var. *angustifolia* T. & G., l. c. (1838) as *G. volubilis*, var. *intermedia* Vail. l. c. 508, changing the name because of an earlier *G. angustifolia* Kunth, Mimos. t. 56 (1824). *G. pilosa*, var. *angustifolia* T. & G. and *G. volubilis*, var. *intermedia* are *G. parvifolia* A. Richard, Essai Fl. Cuba, i. 414 (1845). Should it be felt that this smaller plant of the West Indies and southern Florida, with short racemes and legumes, is only a variety of *G. Macreei* it is clear that the latter name, published in 1837, has precedence over *G. parvifolia* (1845). The varietal name, *G. pilosa*, var. *angustifolia* T. & G. (1838), in no way based upon *G. angustifolia* Kunth (1824), would be the correct one to take up.

When Torrey & Gray changed the spelling of Curtis's species, from near Wilmington, North Carolina, from the original *Macreei* to "*Macraei*", as *G. pilosa* β . *Macraei*, they presumably associated it with the Canadian W. F. Macrae, who sent plants to them, one of which was named for him as *Corallorhiza Macraei* Gray, Gray citing him as "*W. F. Macrae*." In their preface (xiii) Torrey & Gray acknowledged the help of the Canadian "Mr. Macrae." As a matter of fact, Curtis, as indicated on his p. 84, was naming his *Galactia* for a Carolina botanist: "Several [species] are furnished by Dr. McRee, from his plantation, at Rocky Point, a few miles north of Wilming-

ton." In a note on this North Carolina botanist, Barnhart¹ completes the data: JAMES FERGUS McREE (1794–1869), born near Wilmington, M. D. (College of Physic. and Surg., N. Y.), 1814.

**LINUM FLORIDANUM* (Trel.) Planch. DINWIDDIE COUNTY: border of dry sandy woods near Carson, *F. L. & S.*, no. 5817. SUSSEX COUNTY: dry argillaceous field north of Littleton, *F. & L.*, no. 6251.

For discussion see Fernald, *RHODORA*, xxxvii. 429, pl. 396, figs. 11–14 (1935).

**POLYGALA HARPERI* Small. SUSSEX COUNTY: Waverly, 1891, *A. B. Seymour*, no. 6; grassy roadside southeast of Waverly, *F. & L.*, no. 6258; dry pinelands about 4 miles northwest of Waverly, *F. & L.*, no. 6261. See p. 346.

P. RAMOSA Ell. SUSSEX COUNTY: depressions in argillaceous field north of Littleton, *F. & L.*, no. 6263. See p. 339.

STILLINGIA SYLVATICA L. ISLE OF WIGHT COUNTY: "collected in Isle of Wight County, near Franklin," 1893, *Heller*, no. 921; border of dry sandy woods near Joyner's Bridge, *F. G. & L.*, no. 6627. See p. 357.

**AESCLUS DISCOLOR* Pursh. GREENSVILLE COUNTY: large shrub in bottomland woods along Caney Branch, east of Emporia, *F. G. & L.*, no. 6633. See p. 352.

CYRILLA RACEMIFLORA L. SOUTHAMPTON COUNTY: about Franklin, 1893, *Heller*, no. 1032, "new to northern range"; Franklin, 1909, *W. W. Eggleston*, no. 4917; margin of cypress swamp by Blackwater River, near Oak Grove School, *F. & L.*, no. 6630. Several collections from Norfolk County. See p. 359.

HIBISCUS MILITARIS Cav. Seen by us only along the Nottoway River in SOUTHAMPTON COUNTY: Cypress Bridge, *F. & L.*, no. 6274; Courtland, *F. & L.*, no. 6834. See p. 340.

HYPERICUM SETOSUM L. PRINCE GEORGE COUNTY: argillaceous and siliceous boggy depression southeast of Petersburg, at head of Poo Run, *F. L. & S.*, no. 5844, *F. & L.*, no. 6278. NANSEMOND COUNTY: damp sandy and peaty woods and margin of bordering ditch, southwest of Whaleyville, *F. & L.*, no. 6836. See pp. 324, 335 and 364.

**H. DENTICULATUM* Walt., var. *OVALIFOLIUM* (Britton) Blake. SUSSEX COUNTY: sandy and peaty depression (exsiccated shallow pond) about 4 miles northwest of Homeville, *F. & L.*, no. 6837. See p. 337.

**H. DISSIMULATUM* Bickn. PRINCE GEORGE COUNTY: exsiccated argillaceous swale about 3 miles southeast of New Bohemia, *F. G. & L.*, no. 6638. ISLE OF WIGHT COUNTY: sandy roadside ditch south of Zuni, *F. G. & L.*, no. 6639. See p. 353.

**H. PETIOLATUM* Walt., var. *TUBULOSUM* (Walt.) Fern. in *RHODORA*,

¹Jour. N. Y. Bot. Gard. xxi. 167 (1920).

xxxviii. 436 (1936). *Triadenum longifolium* Small. SOUTHAMPTON COUNTY: sandy wooded bottomland of Nottoway River, Courtland, *F. & L.*, no. 6646. See p. 358.

LECHEA.

Pending the publication in the next volume of RHODORA of Dr. Hodgdon's monograph of *Lechea*, the Virginia records are withheld.

*VIOLA AFFINIS Le Conte, var. CHALCHOSPERMA (Brainerd) Griscom. SOUTHAMPTON COUNTY: siliceous and argillaceous alluvium bordering cypress swamp, bottomland of Nottoway River, above Cypress Bridge, *F. & L.*, no. 6289. See p. 341.

*VIOLA LANCEOLATA L., var. VITTATA (Greene) Weath. & Grisc. SUSSEX COUNTY: sandy and peaty depression (exsiccated shallow pond), about 4 miles northwest of Homeville, *F. & L.*, no. 6290. See p. 337.

*LYTHRUM LANCEOLATUM Ell. SUSSEX COUNTY: wet sandy thicket, Burt, *F. & L.*, no. 6295. See p. 342.

RHEXIA VENTRICOSA Fern. & Grisc. PRINCE GEORGE COUNTY: fallow argillaceous field east of Prince George, *F. L. & S.*, no. 5855. CHESTERFIELD COUNTY: exsiccated argillaceous swale west of Petersburg Turnpike, north of Swift Creek, *F. & L.*, no. 6301. See p. 344.

R. MARIANA L., var. PURPUREA Michx. PRINCE GEORGE COUNTY: sphagnous boggy swale southeast of Petersburg, at head of Poo Run, *F. & L.*, no. 6300.

Extension from Southampton and Norfolk Counties.

R. CILIOSA Michx. PRINCE GEORGE COUNTY: dryish upper border of sphagnous boggy swale, about 3 miles southeast of Petersburg, at head of Poo Run, *F. & L.*, nos. 6296 and 6626. ISLE OF WIGHT COUNTY: sphagnous depression in sandy pine woods south of Zuni, *F. & L.*, no. 6625. NANSEMOND COUNTY: damp sandy and peaty woods and margin of bordering ditch, southwest of Whaleyville, *F. & L.*, no. 6841. See pp. 335, 358 and 364.

LUDWIGIA HIRTELLA Raf. Locally abundant in peaty depressions and boggy swales of CHESTERFIELD, PRINCE GEORGE, SUSSEX and NANSEMOND COUNTIES. See p. 335.

L. LINEARIS Walt. PRINCE GEORGE COUNTY: sphagnous boggy swale, about 3 miles southeast of Petersburg, at head of Poo Run, *F. & L.*, no. 6652. SOUTHAMPTON COUNTY: sandy wooded swamp southwest of Cypress Bridge, *F. & L.*, no. 6306. See p. 335.

*OENOTHERA FRUTICOSA L., var. HUMIFUSA T. F. Allen. SUSSEX COUNTY: fallow ploughed field in pineland, about 4 miles northwest of Waverly, *F. & L.*, no. 6440, *F. G. & L.*, no. 6657. See p. 347.

*OE. FRUTICOSA, var. EAMESII (Robinson) Blake. SUSSEX COUNTY: with the preceding, *F. & L.*, no. 6439, *F. G. & L.*, no. 6658. See p. 347.

PROSERPINACA PECTINATA Lam. ISLE OF WIGHT COUNTY: swampy depressions in sandy woods, south of Zuni, *F. G. & L.*, no. 6660. See p. 354.



Photo. E. C. Ogden.

LYSIMACHIA LANCEOLATA: FIG. 1, base, $\times 1$, from North Carolina; FIG. 2, calyx, $\times 4$, from Virginia; FIG. 3, bases of cauline leaves, $\times 4$, from North Carolina; FIG. 4, rosette-leaf, $\times 4$, from Pennsylvania.

L. HYBRIDA: FIG. 5, base, $\times 1$, from Massachusetts; FIG. 6, bases of cauline-leaves, $\times 4$, from New Jersey; FIG. 7, calyx, $\times 4$, from New Jersey.




Photo. H. G. Fernald.

GALIUM CIRCAEZANS, var. *TYPICUM*: FIG. 1, one of the *TYPE* specimens, $\times \frac{1}{2}$, courtesy of Professor H. HUMBERT; FIG. 2, lower leaf-surface, $\times 5$, of plant from Georgia.

Var. *HYPOMALACUM*: FIG. 3, *TYPE*, $\times \frac{1}{2}$; FIG. 4, lower leaf-surface, $\times 5$.

SANICULA GREGARIA Bickn. CHESTERFIELD COUNTY: wooded river-swamp along Appomattox River, near Hopewell, *F. L. & S.*, no. 5866.

*S. CANADENSIS L., var. FLORIDANA (Bickn.) H. Wolff. ISLE OF WIGHT COUNTY: dry sandy yellow pine and oak woods north of Walters, *F. G. & L.*, no. 6661. See p. 354.

HYDROCOTYLE CANBYI C. & R. ISLE OF WIGHT COUNTY: along ditch bordering swampy woods, east of Joyner's Bridge, *F. & L.*, no. 6846.

Extension inland from Princess Anne County.

H. RANUNCULOIDES L.f. SURRY COUNTY: margin of pond in cypress swamp, Sunken Meadow Beach, *F. & L.*, no. 6845. SUSSEX COUNTY: pool in pinelands about 4 miles northwest of Waverly, *F. & L.*, no. 6314.

Extension inland from Princess Anne County.

OXYPOLIS RIGIDIOR (L.) C. & R. PRINCE GEORGE COUNTY: argillaceous and siliceous boggy depressions, about 3 miles southeast of Petersburg, at head of Poo Run, *F. L. & S.*, no. 6848.

The only time seen by us south of the James.

KALMIA ANGUSTIFOLIA L. ISLE OF WIGHT COUNTY: "near Franklin," 1893, *Heller*, no. 1124; dry sandy woods and pine barrens south of Zuni, *F. G. & L.*, nos. 6665 and 6666. See p. 354.

The only area in the southeastern counties in which we have seen this essentially northern (even Hudsonian) species. The Isle of Wight material is transitional to the glandless southern extreme:

K. ANGUSTIFOLIA L., var. **carolina** (Small), comb. nov. *K. carolina* Small, Fl. Se. U. S. 886, 1336 (1903). The only Virginia material of the variety seen by me is from region of Dismal Swamp, *Biltmore Herb.*, no. 1344^c.

LYONIA LUCIDA (Lam.) C. Koch. SOUTHAMPTON COUNTY: margin of cypress swamp by Blackwater River, near Oak Grove School, *F. & L.*, no. 6667. See p. 359.

*GAYLUSSACIA DUMOSA (Andr.) Torr. & Gray, var. BIGELOVIANA Fernald. PRINCE GEORGE COUNTY: border of dry woods east of PRINCE GEORGE, *F. L. & S.*, no. 5874.

The copiously glandular northern extreme. In the pine-barren areas only typical southern *G. dumosa* was seen.

VACCINIUM ELLIOTII Chapm. Tall, often tree-like shrub up to 3.5 m. high, abundant in thickets and along streams in SOUTHAMPTON COUNTY. See p. 331.

*PYXIDANTHERA BARBULATA Michx. ISLE OF WIGHT COUNTY: dry sandy pine barrens, south of Zuni, *F. G. & L.*, no. 6669, *F. & L.*, no. 7137. See p. 355.

GALAX APHYLLA L. To the few Coastal Plain stations add ISLE OF WIGHT COUNTY: rich wooded bank of Blackwater River near Joyner's Bridge, *F. G. & L.*, no. 6668.

LYSIMACHIA RADICANS Hook. (*Steironema radicans* (Hook.) Gray.) SOUTHAMPTON COUNTY: siliceous and argillaceous alluvium bordering cypress swamp, bottomland of Nottoway River, above Cypress Bridge, *F. & L.*, no. 6332, distributed as *L. lanceolata*, var. *hybrida*. See p. 341.

Lysimachia radicans, characterized by its sprawling or arching habit, with the prolonged stems rooting at nodes and soon reclining, with long-petioled and membranaceous lanceolate to lance-ovate blades, and with small nodding flowers 8–12 mm. long, the calyx-lobes only 3–5 mm. long and exceeded by the capsule, has been standing in our manuals as a plant of Virginia. In the Gray Herbarium *L. radicans* has heretofore been represented only from the Mississippi drainage, Mississippi to eastern Texas, northward to western Tennessee and Missouri. Our material, though very inadequate, can be matched only in that species, but it shows an inclination to whorled leaves and its flowers are all solitary on simple peduncles in the axils of the primary leaves, instead of being borne on axillary branches as in most true *L. radicans*. Fuller material may show the plant of southeastern Virginia to be distinct.

THE IDENTITY OF *LYSIMACHIA LANCEOLATA* (PLATE 482). In preparing the Synoptical Flora of North America Asa Gray revived the genus *Steironema* Raf. for an American group, which some later authors, for instance Knuth¹ and Handel-Mazzetti,² treating the genus from a world-viewpoint, have put back as a section or a subsection into *Lysimachia*. Handel-Mazzetti shows that the characters relied upon to keep American *Steironema* apart break down in some Chinese species of *Lysimachia*, and that *Steironema* "cannot be treated as a different genus as has been done recently by some American botanists." Returning *Steironema* to *Lysimachia* is simple enough; the difficulty is in identification of some of the older types.

When Gray reviewed the plants which he treated as *Steironema lanceolatum* (Walt.) Gray, based upon *Lysimachia lanceolata* Walt. Fl. Carol. 92 (1788), he made it an inclusive species without clearly defined varieties. He published the combinations in 1876, in Proc.

¹ Knuth in Engler, *Pflanzenr.* iv²³⁷, 257, 276 (1905).

² Handel-Mazzetti in *Notes, Roy. Bot. Gard. Edinb.* xvi. 52 (1928) and in *Die Pflanzenareale*, 2 Reihe, v. Karten 44–49 (1929).

Am. Acad. xii. 63, but his definitions were published in the Synoptical Flora:

S. lanceolatum, Gray. Stems erect, a foot or two high, simple or paniculately branched, somewhat angled: leaves lanceolate or linear, an inch or two long, tapering into a short and margined ciliate petiole or attenuated base; the radical and sometimes lowest cauline from oblong to orbicular, small: corolla about two thirds inch in diameter; its divisions conspicuously erose and cuspidate-acuminate, slightly exceeding the lanceolate calyx-lobes.—Proc. Am. Acad. l. c. *S. heterophylla*, Raf. l. c. *S. florida*, Baudo, l. c., chiefly. *Anagallis lutea*, &c., Pluk. Alm. t. 333, f. 1. *Lysimachia lanceolata*, Walt. Car. 92. *L. hybrida* & *heterophylla*, Michx. Fl. i. 126. *L. ciliata*, var., Chapm. Fl. 280. *L. decipiens*, Bertoloni, Amoen.—Low grounds and thickets, western parts of Canada to Florida, and Nebraska to Louisiana. Polymorphous; the extremes in the following varieties, the first of which verges to the two preceding species.

Var. hybridum. Cauline leaves mostly petioled, from oblong to broadly linear.—*Lysimachia lanceolata*, var. *hybrida*, Gray, l. c. *L. hybrida*, Michx. l. c. *L. heterophylla*, Ell., Nutt., &c.—Commoner northward and westward.

Var. angustifolium. Stems more branched, a span to 2 feet high: cauline leaves linear, acute at both ends, more sessile, a line or two broad.—*L. angustifolia*, Lam. Ill. i. 440, not Michx. *L. heterophylla*, Michx. l. c. *L. quadriflora*, Ell., hardly of Bot. Mag.—The more marked form mainly southward.¹

In his earlier paper Gray had noted that "The species are not easy to define, as they incline to run into each other." This attitude toward them may account for Gray's inclusion of *Lysimachia heterophylla* Michx. Fl. Bor.-Am. i. 126 (1803) in the synonymy of both his *Steironema lanceolatum* (typical) and his var. *angustifolium* (Lam.) Gray, which rested upon *L. angustifolium* Lam. Ill. i. 440 (1797?). Subsequent authors, leaving *Steironema lanceolatum* much as defined by Gray, have separated *S. heterophyllum* (Michx.) Raf. from it as a species. There are certainly two well defined species included in Gray's general concept. In dry to moist open woods or thickets or in swales and on shores, but mostly in dryish habitats in the South, from Florida to Louisiana, northward into Pennsylvania, Ohio, southern Michigan and Wisconsin, is a species which for the time being may be called

No. 1. Stems slender and firm, 0.5–7 dm. high, from elongate cord-like or filiform rhizomes and stolons, simple or with ascending branches (the latter often abbreviated); basal leaves often rosulate, oblong, elliptic or rounded, petioled; middle and upper leaves linear to lanceolate or narrowly oblong, bristly-ciliate at base, sessile or subsessile,

¹ Gray, Synop. Fl. N. Am. ii. 61, 62 (1878).

pale beneath; calyx-segments firm, their lateral nerves not evident. FIGS. 1-4.

Farther north, extending from Quebec to western Ontario and North Dakota, southward through the northeastern states and more locally to Florida and Texas, is a coarser plant of wet shores, sloughs and swamps. This may be called

No. 2. Stems stoutish, from a soft base, without stolons or slender rhizomes, 0.2-1.5 m. long, ascending or, when very elongate, becoming procumbent, the autumnal basal rosettes sessile or on short thick offshoots; cauline leaves linear-lanceolate to oblong, mostly petioled, green on both sides, the petiole, but rarely the blade, somewhat ciliate; calyx-segments herbaceous, 3-nerved. FIGS. 5-7.

No. 1 is the plant described very clearly, though briefly, by Michx. (1803) as *Lysimachia*:

HETEROPHYLLA. *L. gracilis*, glabra: foliis oppositis; imis suborbiculatis et brevi-petiolatis; superioribus linearibus, sessilibus, basi ciliolatis: floribus cernuis.

Obs. Flores omnino *LYSIMACHIAE ciliatae*.

PLUCK. *mantiss.* t. 333. fig. I. Affinis.

Hab. in Georgia.¹

It had with almost equal clarity been defined some years earlier by Lamarck as

1977 *LYSIMACHIA angustifolia*.

L. foliis linearibus, basi ciliatis, sessilibus; pedunculis unifloris; corollis calyce brevioribus.

E Carolina. *D. Fraser*.²

These, it will be noted, were both included by Gray under his *Steironema lanceolatum*, var. *angustifolium*, although *L. heterophylla* Michx. was also put by him under typical *S. lanceolatum*.

Both nos. 1 and 2 may have leaves of any outline from linear or linear-lanceolate and very narrow through broader-lanceolate to oblong. It becomes evident that Gray, not cognizant of the very different bases and other characters of nos. 1 and 2, was merely putting plants with "leaves lanceolate or linear, . . . tapering into a short . . . petiole" into his *Steironema lanceolatum*; those with "cauline leaves linear, . . . more sessile, a line or two broad" into his var. *angustifolium*; and those with "Cauline leaves mostly petioled, from oblong to broadly linear" into his var. *hybridum*, based on *Lysimachia hybrida* Michx. The really diagnostic characters were

¹ Michx. Fl. Bor.-Am. i. 127 (1803).

² Lam. Ill. i. 440 (1797 or earlier).

not noted by him. It consequently becomes significant, in going back to Walter's original account of his *L. lanceolata* to read:

lanceolata foliis lanceolatis subsessilibus, petalis acumine terminatis.¹
2.

The subsessile leaves and the abundance of our no. 1 (*L. angustifolia* Lam. and *L. heterophylla* Michx.) in Walter's territory and the rarity there (if it occurs at all) of our no. 2 make Gray's own examination of the Walter type of utmost importance. Studying Walter's herbarium on February 9th, 1839, Gray, with more modern American specimens for comparison, made the memorandum: "*Lysimachia lanceolata!* = mine from Michigan." This Michigan specimen, ticketed "HERB. A. GRAY" and marked in Gray's hand: "Michigan State Coll.", is very typical broad-leaved *L. angustifolia* Lam. or *L. heterophylla* Michx., showing clearly the cord-like rhizome, the round-tipped basal leaves, the subsessile ciliate-based cauline ones with the characteristic grayish sheen beneath, and the firm sepals. This plant, positively identified by Gray in 1839 with Walter's type, seems to settle the identity of *L. lanceolata* Walt. Singularly enough, however, in the Synoptical Flora, where he relied primarily on leaf-outline, Gray so far forgot his comparison of 1839 as to ticket the Michigan specimen as "*Steironema lanceolatum*, var. *hybridum*," thus making the already confounded confusion still worse!

With no. 1 of page 439 reasonably settled as *LYSIMACHIA LANCEOLATA* Walt., the proper name must be found for no. 2, the coarser, thick-stemmed nonstoloniferous plant with middle and upper leaves more petioled and green beneath, and with herbaceous 3-nerved calyx-lobes. Apparently the oldest name for it is *L. HYBRIDA* Michx. Fl. Bor.-Am. i. 126 (1803). There is no reasonable doubt of the identity. Michaux was distinguishing our no. 1 as his *L. heterophylla*. His *L. hybrida* has "foliis oppositis, longe petiolatis, lanceolatis, basi sensim acutis; petiolo ciliato," etc.; and when I examined it in 1903, familiar only with the coarse plant of New England, I made the note: "*hybrida*. The common *lance*-leaved plant."

Some of the more recent names of Rafinesque, Greene and others doubtless designate variations in leaf-outline of the heteromorphic *Lysimachia lanceolata* and *L. hybrida*. I leave their interpretation to those who see value in them. One of Greene's proposed species of the

¹ Walt. Fl. Carol. 92 (1788).

group is *Steironema pumilum*. In Rydberg's Flora of the Prairies and Plains is the following key:

"Leaf-blades mostly ovate or ovate-lanceolate, rounded or subcordate at the base; 1. *S. pumilum*.
Leaf-blades mostly lanceolate, cuneate at the base; 2. *S. ciliatum*."

Nevertheless, *S. ciliatum* is thus described just below: "blades ovate or lanceolate, . . . , acute, rounded, truncate, or subcordate at the base."

LYSIMACHIA LANCEOLATA is local in southeastern Virginia. JAMES CITY COUNTY: margin of dried-up pond $\frac{1}{2}$ mile south of Ewell, Grimes, no. 4481. HENRICO COUNTY: exsiccated argillaceous swale, Libbie Avenue, Westhampton, *F. L. & S.*, no. 5888, distributed as *L. heterophylla*. SUSSEX COUNTY: rich oak woods near Moore's Mill, *F. & L.*, no. 7139.

*FRAXINUS PROFUNDA Bush. NORFOLK COUNTY: gum swamps and wet woods near Indian Creek, *F. & G.*, no. 4690, distributed as *F. pennsylvanica*. SOUTHAMPTON COUNTY: sandy alluvial bottomlands of Three Creek, Drewryville, *F. L. & S.*, no. 5891. See p. 331.

Heretofore known from Louisiana to Georgia, north in the lower regions to Illinois, Indiana and Ohio.

FRAXINUS CAROLINIANA Mill., var. **pubescens** (M. A. Curtis), comb. nov. *F. platycarpa*, β . *pubescens* M. A. Curtis in Am. Journ. Sci. ser. 2, vii. 408 (1849). *F. Rehderiana* Lingelsheim in Engler, Pflanzenr. iv²⁴³. 42 (1920). *F. caroliniana*, var. *Rehderiana* (Lingels.), Sargent in Journ. Arn. Arb. ii. 173 (1921).—Quite as common as the glabrous-leaved typical *F. caroliniana*.

*LIGUSTRUM SINENSE Lour. YORK COUNTY: border of dry woods, 2 miles south of Yorktown, *F. L. & F.*, no. 4991. NANSEMOND COUNTY: dry sandy woods and adjacent clearings, *F. L. & F.*, no. 4990. ISLE OF WIGHT COUNTY: border of dry sandy woods south of Zuni, *F. G. & L.*, no. 6670. See p. 353.

CYNOCTONUM MITREOLA (L.) Britton. PRINCE GEORGE COUNTY: exsiccated argillaceous swale about 3 miles southeast of New Bohemia, *F. & L.*, no. 6339. ISLE OF WIGHT COUNTY: muddy margin of Blackwater River, near Joyner's Bridge, *F. & L.*, no. 6851. See pp. 345, 346 and 362.

SABATIA PANICULATA (Michx.) Pursh. Frequent in dry argillaceous fields, thickets and clearings of CHESTERFIELD and SUSSEX COUNTIES. See p. 339.

The milk-white corollas always change to yellowish or saffron-colored in the herbarium. Thinking that quick drying might save the color, we employed this method, but specimens which seemed

satisfactory when they came from press had lost their whiteness in a few weeks.

S. BRACHIATA Ell. SUSSEX COUNTY: sandy woods and clearings northwest of Homeville, *F. & L.*, no. 6344; dry argillaceous field north of Littleton, *F. & L.*, no. 6345; seen in abundance near Waverly. See p. 339.

Typical *Sabatia brachiata* has very handsome rosy-pink corollas. Occasional albinos are found, which in the field (see p. 339) strongly suggest *S. paniculata*. Their corollas however, are larger and they do not change to saffron-color after drying. The albino may be called

S. BRACHIATA*, forma **candida, f. nov., corollis albidis.—VIRGINIA: Waverly, July 20, 1891, *A. B. Seymour*, no. 33, as *S. paniculata*; dry argillaceous field north of Littleton, July 22, 1936, *Fernald & Long*, no. 6346 (TYPE in Gray Herb.). See p. 339.

S. CALYCINA (Lam.) Heller. SOUTHAMPTON COUNTY: in swampy ground, Franklin, 1893, *Heller*, no. 1114¹; border of muddy pool in Three Creek, Drewryville, *F. L. & S.*, no. 5895; siliceous and argillaceous alluvium bordering cypress swamp, bottomland of Nottoway River, above Cypress Bridge, *F. & L.*, no. 6348. ISLE OF WIGHT COUNTY: sandy alluvial woods, bottomland of Blackwater River, Zuni, *F. & L.*, no. 6349. See p. 340.

S. CAMPANULATA (L.) Torr. PRINCE GEORGE COUNTY: argillaceous and siliceous boggy depression southeast of Petersburg, at head of Poo Run, *F. & L.*, no. 6350. SUSSEX COUNTY: sandy and peaty depression (exsiccated shallow pond), about 4 miles northeast of Homeville, *F. & L.*, no. 6351. See p. 337.

Typical *Sabatia campanulata* or *Chironia campanulata* L. (1753), upon which it rests, is, as shown by a photograph supplied by Mr. Savage, the plant which ranges from Massachusetts to eastern Virginia, thence taking to the mountains of North and South Carolina and southward to southern Georgia and Alabama. In this plant the primary cauline leaves are oblong-linear to lanceolate, the pedicels are naked or only slightly bracted, the linear calyx-segments (except in small secondary flowers) 1–2 cm. long, the corolla-segments 1–1.7 cm. long. On the Coastal Plain from Florida to Louisiana and North Carolina occurs *S. gracilis* Michx., which is commonly reduced outright to *S. campanulata*. In its best development, however, it is smaller throughout, with the lower cauline leaves linear, the upper very narrowly so, the pedicels mostly leafy-bracted, the linear-acicular calyx-segments 6–14 mm. long, the corolla-segments 6–14 mm. long. The material from southeastern Virginia stands midway

¹ See Heller, l. c. 24 (1894).

between most typical *S. campanulata* and *S. gracilis*, having the narrow leaves and calyx-segments of the latter but many of the pedicels naked, and the long calyx and large corolla of the former. This transitional series in southeastern Virginia makes it clear that *S. gracilis* should be treated as a geographic variety:

SABATIA CAMPANULATA (L.) Torr., var. **gracilis** (Michx.), comb. nov. *Chironia gracilis* Michx. Fl. Bor.-Am. i. 146 (1803).

GENTIANA PORPHYRIO J. F. Gmel. NANSEMOND COUNTY: very rare at border of dry sandy pine woods south of Factory Hill, *F. & L.*, no. 6852. See p. 364.

BARTONIA PANICULATA (Michx.) Muhl. PRINCE GEORGE COUNTY: argillaceous and siliceous boggy depressions, about 3 miles southeast of Petersburg, at head of Poo Run, *F. L. & S.*, no. 6860. NANSEMOND COUNTY: damp sandy and peaty woods and margin of bordering ditch, southwest of Whaleyville, *F. & L.*, no. 6859.

TRACHYLOSPERMUM DIFFORME (Walt.) Gray. Frequent in damp thickets and at borders of wet woods, northward to HENRICO COUNTY: exsiccated argillaceous swale, Libbie Avenue, Westhampton, *F. L. & S.*, no. 5897.

Here noted because not included in Merriman's *Flora of Richmond and Vicinity*.

ASCLEPIAS RUBRA L. PRINCE GEORGE COUNTY: argillaceous and siliceous boggy depressions, about 3 miles southeast of Petersburg, at head of Poo Run, *F. L. & S.*, no. 5901; similar habitat on headwaters of Blackwater River, *F. L. & S.*, no. 5902. See p. 326.

ACERATES VIRIDIFLORA (Raf.) Eaton. PRINCE GEORGE COUNTY: dry pineland west of Prince George, *F. & L.*, no. 5900.

The only time seen by us in the southeastern counties.

BREWERIA HUMISTRATA (Walt.) Gray. Frequent in dry sandy woods and openings, SOUTHAMPTON and SUSSEX COUNTIES. See pp. 333 and 339.

HYDROLEA QUADRIVALVIS Walt. SUSSEX COUNTY: water-hole in sandy and peaty depression (exsiccated shallow pond), about 4 miles northwest of Homeville, *F. & L.*, nos. 6362 and 6671. SOUTHAMPTON COUNTY: sandy alluvial bottomlands of Three Creek, Drewryville, *F. L. & S.*, no. 5864. See pp. 332 and 338.

Although the species appears in manuals as a native of Virginia, our collections seem to be the first from north of the extreme southern boundary of the state. Heller's collection (no. 1162) of 1893, the previous basis, is slightly equivocal. It bears a label, headed "Plants of Northeastern North Carolina. Collected near Margarettsville, Northampton Co.," but the label bears the annotation "On the Va. line."¹

¹ See Heller. l. c. 25 (1894).



Photo. E. C. Ogden.

EUPATORIUM ALBUM, var. TYPICUM: FIG. 1, leaf, $\times 2$, from Virginia; FIG. 2, involucre, $\times 4$, from New Jersey.

Var. GLANDULOSUM: FIG. 3, involucre, $\times 4$, from South Carolina.

Var. MONARDIFOLIUM: FIG. 4, portion of TYPE, $\times \frac{2}{5}$; FIG. 5, leaf, $\times 2$, from North Carolina; FIG. 6, involucre, $\times 4$, from TYPE.

Var. SUBVENOSUM: FIG. 7, leaf, $\times 2$, from TYPE; FIG. 8, involucre, $\times 4$, from TYPE.



Photo. E. C. Ogden.

EUPATORIUM LEUCOLEPIS: FIG. 1, characteristic foliage, $\times \frac{2}{5}$, from New Jersey; FIG. 2, stem and leaf-bases, $\times 4$, from Virginia.

Var. NOVAE-ANGLIAE: FIG. 3, type, $\times \frac{2}{5}$; FIG. 4, stem and base of median leaf, $\times 4$; FIG. 5, stem and upper leaves, $\times 4$.

HELIOTROPIUM EUROPAEUM L. HENRICO COUNTY: waste places and roadsides, Richmond, *F. L. & S.*, no. 5904.

H. INDICUM L. SOUTHAMPTON COUNTY: sandy alluvial bottomland of Nottoway River, Courtland, *F. & L.*, no. 6862. See p. 366.

ONOSMODIUM VIRGINIANUM (L.) A. DC. DINWIDDIE COUNTY: border of dry sandy woods near Carson, *F. L. & S.*, no. 5905.

The only time seen in the southeastern counties.

*VERBENA SCABRA Vahl. SURRY COUNTY: border of tidal marsh along Gray's Creek, near Cross Creek Landing, south of Swann Point, *F. & L.*, no. 6863. See p. 365.

*TRICHOSTEMA DICHOTOMUM L., var. **puberulum** Fernald & Griscom, var. nov., caulibus puberulis, ramis similibus vel minute glandulosis vix pilosis.—Mississippi to Florida, north locally to southeastern Virginia. TYPE: Duval County, Florida, *Curtiss*, no. 1976 (in Gray Herb.).

The only material we have seen from north of Florida is our collection from Cedar Hill, NORFOLK COUNTY, no. 2884. All material seen from Florida and Mississippi is clearly of this variety.

Typical *T. dichotomum* L., as ascertained by Mr. C. A. Weatherby on studying the type and as shown by a photograph of it sent from the Linnean Society, is the common and wide-spread plant which has much longer (pilose) pubescence and abundant, usually longer-stalked, glands. Transitional material is common from southern Cape May, New Jersey to southeastern Virginia.

T. LINEARE Walt. ISLE OF WIGHT COUNTY: dry sandy yellow pine and oak woods north of Walters, *F. G. & L.*, no. 6675; border of dry sandy woods near Joyner's Bridge, *F. G. & L.*, no. 6676. See p. 357.

PHYSOSTEGIA DENTICULATA (Ait.) Britton. SOUTHAMPTON COUNTY: siliceous and argillaceous alluvium bordering cypress swamp, bottomland of Nottoway River, above Cypress Bridge, *F. & L.*, no. 6368. ISLE OF WIGHT COUNTY: sandy alluvial woods, bottomland of Blackwater River, Zuni, *F. & L.*, no. 6369. See p. 340.

PYCNANTHEMUM CLINOPODIOIDES T. & G., NORFOLK COUNTY: border of rich woods, south of Great Bridge, *F. L. & F.*, no. 5021. NANSEMOND COUNTY: dry sandy woods and adjacent clearings, Kilby, *F. L. & F.*, nos. 5023–5025. SUSSEX COUNTY: sandy woods and clearings northeast of Homeville, *F. & L.*, no. 6380.

An inland species rare on the Coastal Plain.

*PYCNANTHEMUM PYCNANTHEMOIDES (Leavenworth) Fernald, var. **viridifolium**, var. nov., foliis primariis subtus viridescentibus vix canescentibus hispida; calycis dentibus apice sparse setosis.—Greensville County, VIRGINIA: in clay at border of a dry thicket, near James River Junction, August 19, 1936, *Fernald, Griscom & Long*, no. 6678

(TYPE in Gray Herb.; ISOTYPES in Herb. Phil. Acad., Herb. Griseb. and elsewhere). See p. 353.

Typical *Pycnanthemum pycnanthemoides* of the mountains from Virginia and Kentucky to Georgia and Alabama, has, as originally described by Leavenworth (as *Tullia pycnanthemoides* from eastern Tennessee) the leaves whitened beneath. Their lower surfaces are canescent with minute soft pubescence and the calyx-lobes are abundantly supplied with setae. Var. *viridifolium* is a coastal-plain extreme, with the large oval leaves and the very large lilac-purple and conspicuously spotted corolla of the mountain plant, but only the uppermost or bracteal leaves are whitened, the others green and rather coarsely hispid beneath, while the calyx-teeth have only a single (rarely more) terminal bristle. Exactly the same variation is represented in the Gray Herbarium by an old specimen (without locality but presumably near Santee Canal) from *H. W. Ravenel*; and one of Asa Gray's collections (again without stated locality) shows mixed with more characteristic *P. pycnanthemoides* from "Mts. Carol. 1843," a specimen with the green leaves and the pubescence of var. *viridifolium* but with the more bristly calyx-teeth of the mountain plant.¹

P. VIRGINIANUM (L.) Durand & Jackson. SUSSEX COUNTY: wet sandy thicket, Burt, *F. & L.*, no. 6376.

The only time seen in the southeastern counties.

LYCOPUS EUROPAEUS L. SURRY COUNTY: roadside by sandy thicket, Sunken Meadow Beach, *F. & L.*, no. 6865.

**L. AMERICANUS* Muhl., var. *LONGII* Benner. PRINCE GEORGE COUNTY: argillaceous and siliceous boggy depressions, about 3 miles southeast of Petersburg, at head of Poo Run, *F. L. & S.*, no. 5910. SUSSEX COUNTY: water-hole in sandy and peaty depression (exsiccated shallow pond), about 4 miles northwest of Homeville, *F. & L.*, nos. 6382 and 6681. NANSEMOND COUNTY: ditches bordering sandy woods, Factory Hill, *F. & L.*, no. 6682. See p. 329.

**MICRANTHEMUM UMBROSUM* (Walt.) Blake. SOUTHAMPTON COUNTY: sandy alluvial bottomlands of Three Creek, Drewryville, *F. L. & S.*, no. 5913. ISLE OF WIGHT COUNTY: muddy margin of Blackwater River, near Joyner's Bridge, *F. & L.*, no. 6866. See pp. 331 and 362 and MAP 8.

CHELONE CUTHBERTII Small. PRINCE GEORGE COUNTY: swampy woods west of New Bohemia, *F. L. & S.*, no. 5911. SOUTHAMPTON

¹ In studying this series it has been found necessary to make the following combination:

PYCNANTHEMUM Beadlei (Small), comb. nov. *Koellia Beadlei* Small in Bull. Torr. Bot. Cl. xxv. 470 (1898).

COUNTY: sandy wooded swamp southwest of Cypress Bridge, *F. & L.*, no. 6388. See p. 328 and MAP 3.

PENSTEMON AUSTRALIS Small. SOUTHAMPTON COUNTY: dry sandy oak and pine woods northeast of Cypress Bridge, *F. & L.*, no. 6384; sandy thickets and openings along Nottoway River at Sycamore Bend, *F. & L.*, no. 6386. SUSSEX COUNTY: dry sandy hickory and oak woods, Burt, *F. & L.*, no. 6385. ISLE OF WIGHT COUNTY: dry sandy yellow pine and oak woods near Walters, *F. & L.*, no. 6387. See p. 339.

Slight northern extension.

SCROPHULARIA MARILANDICA L. SUSSEX COUNTY: dry sandy hickory and oak woods, Burt, *F. & L.*, no. 6383. See p. 342.

The only time seen in the southeastern counties.

*SEYMERIA CASSIODES (Walt.) Blake. GREENSVILLE COUNTY: sandy clearing north of Emporia, *F. G. & L.*, no. 6690. ISLE OF WIGHT COUNTY: dry sandy pine barrens south of Zuni, *F. G. & L.*, no. 6691, *F. & L.*, no. 6867. See pp. 352 and 355 and MAP 24.

BUCHNERA AMERICANA L. PRINCE GEORGE COUNTY: dry sandy pine woods about 3 miles southeast of Petersburg, on headwaters of Blackwater River, *F. L. & S.*, no. 5918, *F. & L.*, no. 6736.

Apparently the first known station on the Coastal Plain of Virginia, unless Clayton, whose explorations extended west to the Shenandoah Valley, got it in the eastern counties. See p. 329.

SCHWALBEA AMERICANA L. GREENSVILLE COUNTY: sandy clearing north of Emporia, *F. G. & L.*, no. 6695. See p. 352 and MAP 23.

Our material is very mature, much of the fruit already disintegrating, but on the lower internodes and on the intact fruiting calyces the characteristic reflexed hairs are apparent and the anterior sepals are blunt, as reputed in the northeastern series of specimens, which Pennell treats as typical *Schwalbea americana*. In his detailed discussion of the group Pennell cites *S. americana* as occurring from "Massachusetts and New York to Maryland and perhaps Virginia"—Pennell, *Scroph. E. N. Am.* 486 (1935). He there states that *S. americana* was "Based wholly upon Gronovius' *Flora Virginica* 71. 1743, where the plant is described and typified by Clayton's number 33 from Virginia. This, seen in Herb. British Museum (Natural History) at London, is the northern species now considered. How Clayton obtained the plant, and even whether it came from Virginia is uncertain, since Gronovius' label quotes this remark of his: 'A plant very uncommon, wholly unknown to me: though I think it agrees in most respects with the *Clandestina* of Tournefort,' while another slip states: 'Mr. Clayton of Virginia An. 1734. Numb. 33', doubtless recording the year of receipt by Gronovius."

The statement of Clayton that the plant was "wholly unknown to me" apparently simply meant that he did not recognize it, not that he had received it from some source outside Virginia; and his statement hardly justifies doubt of his having personally collected the plant. At least, we now know *Schwalbea americana* from Virginia, whence it was described, although it is possible that Clayton secured it farther to the north in the state, our station being in one of the southernmost counties (bordering North Carolina). From Clayton's map of Virginia, however, it is evident that he was cognizant of the region southward, he showing the Nottoway River (his no. 48) with its tributary, the Blackwater (no. "49 Nigra aqua").

Pennell separates the more southern material (North Carolina and Kentucky to Florida and Louisiana) as *Schwalbea australis* Pennell in Proc. Acad. Nat. Sci. Phila. lxxi. 289 (1920). His key-differences, in his later treatment, are as follows:

- "A. Pubescence of stem, pedicels, and calyx consisting of up-curved, usually shorter hairs; leaf-blades elliptic-oval, usually about 1.5 cm. wide, usually more obscurely veined; anterior calyx-lobes acute or acutish. 1. *S. australis*.
 AA. Pubescence of stem, pedicels, and calyx consisting of recurved hairs; leaf-blades elliptic-lanceolate, rarely over 1 cm. wide, usually evidently veined; anterior calyx-lobes obtuse. 2. *S. americana*."

The more extreme specimens are well marked but minute "up-curved" pubescence occurs on some Massachusetts specimens; in fact, a fine specimen in the Gray Herbarium from Sandwich, Massachusetts, has such pubescence and Pennell has annotated it in the herbarium as *Schwalbea australis*. Its anterior calyx-lobes, however, are blunt as in *S. americana*, but its oval leaves are 2 cm. broad, extreme even for *S. australis* and surely for *S. americana* as defined. Although in the Gray Herbarium Pennell correctly marked this embarrassing plant as *S. australis*, he cites it as *S. americana*. I do not see how it and Curtiss, no. 6742, TYPE of *S. australis*, essentially differ; the latter, as represented in the Gray Herbarium, likewise has its lower leaves 2 cm. broad. Narrower-leaved plants do occur in both the North and the South but of the 7 southern sheets (*S. australis*) 4 have their broadest leaves only 8-13 mm. broad, while the majority of northern plants show their broadest leaves 7-12 mm. broad. I am unable to separate two varieties and much less two species.

PEDICULARIS LANCEOLATA Michx. CHESTERFIELD COUNTY: wooded

river-swamp along Appomattox River near Hopewell, *F. L. & S.*, no. 5919.

Slight extension southward.

[*UTRICULARIA VIRGATULA* Barnhart. In a previous paper, *RHODORA*, xxxviii. 444 (1936), I recorded the station in Northampton County as "An important discovery, giving us a station intermediate between Cape May, New Jersey and Florida and Cuba." Dr. S. F. Blake kindly calls my attention to a station recorded in Hitchcock & Standley's *Flora of the District of Columbia and Vicinity*, 255 (1919) near Suitland, about as far west of Cape May as the Savage Neck station is to the south. I apologize for overlooking the record, somewhat obscured under the alias *Stomoisia virgatula*.]

**CATALPA SPECIOSA* Warder. SOUTHAMPTON COUNTY: border of dry woods, Ivor, probably original trees planted but now well naturalized, *F. & L.*, no. 6396, *F. G. & L.*, no. 6696. See p. 347.

THE TYPE OF *GALIUM CIRCAEZANS* (PLATE 483). *Galium circaezans* consists of two well defined geographic varieties. The wide-ranging plant of the North, from Maine and southwestern Quebec to Minnesota and Nebraska, south in the uplands to interior North Carolina, Kentucky, Missouri and Oklahoma, is coarser than the southern extreme, its larger leaves 2–5 cm. long and 1–2.5 cm. broad, their nerves conspicuously long-hirsute beneath. In the South, from Florida to Texas, the plant is more slender and with smaller and less pubescent leaves, the larger ones only 1.5–2.5 cm. long by 0.7–1.4 cm. broad, the nerves beneath sparingly short-hispid to glabrous. This southern extreme meets the northern in Virginia and elsewhere, extending very locally to Rhode Island, Connecticut, New York, Kentucky and southern Michigan. Impressed by the general occurrence of the small- and smoother-leaved extreme in the South, it has seemed important to determine just what Michaux had before him when he described *G. circaezans* from Carolina. Through the great courtesy of Professor Humbert and the skill of M. Cintract I now have a remarkably clear photograph of the type (FIG. 1). It consists of three fruiting stems of the southern extreme, its blunt, oval leaves 1.5–2 cm. long by 7–12 mm. broad, with very short and scattered hispidity on the veins beneath. It was perfectly described by Michaux, "foliis quaternis, ovalibus, . . . margine ciliato nervisque (armato oculi) hirsutulis"; and it is the blunt-leaved form of the nearly smooth plant described as *G. circaezans* var. *glabrum* Britton, from eastern New York.

Since typical *Galium circaezans* is the chiefly southern extreme with

small and sparsely hispid to glabrous leaves, the more northern extreme may be called

GALIUM CIRCAEZANS Michx., var. **hypomalacum**, var. nov. (TAB. 483, FIG. 3 et 4), foliis majoribus 2–5 cm. longis 1–2.5 cm. latis, nerviis subtus longe hirsutis.—Dry woods, southern Quebec to Minnesota, south to the uplands of North Carolina, Kentucky, Missouri and Oklahoma. TYPE: open dry woods, Peoria, Illinois, July, 1903, *F. E. McDonald* in Gray Herb.

The bibliography of typical *Galium circaezans* follows.

GALIUM CIRCAEZANS Michx., var. **typicum**. *G. circaezans* Michx. Fl. Bor.-Am. i. 80 (1803). *G. boreale?* Walt. Fl. Carol. 87 (1788), not L. *G. circaeoides* R. & S. Syst. iii. 256 (1818). Var. *glabrum* Britton in Bull. Torr. Bot. Cl. xxi. 32 (1894). Var. *glabellum* Britton, Mem. Torr. Bot. Cl. v. 303 (1894).

In plate 483, prepared by Henry G. Fernald, FIG. 1 is one of the TYPE specimens of var. TYPICUM, $\times \frac{1}{2}$; FIG. 2, the lower leaf-surface of var. TYPICUM, $\times 5$, from Marietta, Georgia, *R. N. Larrabee*. FIG. 3 is the TYPE of var. HYPOMALACUM, $\times \frac{1}{2}$; FIG. 4 the lower surface of a leaf, $\times 5$.

HOUSTONIA TENUIFOLIA Nutt. SUSSEX COUNTY: dry sandy hickory and oak woods, Burt, *F. & L.*, no. 6400. See p. 342.

**H. LANCEOLATA* (Poir.) Britton. NANSEMOND COUNTY: Suffolk, July 15, 1895, *J. W. Blankinship*.

A species of the interior of the continent, Alabama to Oklahoma, north to Kentucky, Illinois and Missouri. On the Atlantic slope, locally from eastern Virginia to southern Maine.

Only station seen for this and several companion species on the Coastal Plain.

**OLDENLANDIA BOSCHII* (DC.) Chapm. SOUTHAMPTON COUNTY: open sandy borders of pools and depressions, bottomland of Nottoway River, Courtland, *F. & L.*, no. 6700. See p. 359.

SYMPHORICARPOS ORBICULATUS Moench.

The frequent statement that *Symphoricarpos orbiculatus* is naturalized or a garden escape eastward does not apply to southeastern Virginia. From Dinwiddie to Surry County and southward it is a consistent part of the native thickets and undergrowth. In fact, it was recorded from Virginia by Pursh, in 1814, as *Symphoria glomerata*.

LOBELIA GLANDULIFERA (Gray) Small. PRINCE GEORGE COUNTY: argillaceous and siliceous boggy depressions, about 3 miles southeast of Petersburg, at head of Poo Run, *F. L. & S.*, no. 6877; exsiccated argillaceous swale about 3 miles southeast of New Bohemia, *F. & L.*, nos. 6406 and 6876. See pp. 339, 345 and 363 and MAP 20.

THE VARIETIES OF EUPATORIUM ALBUM (PLATE 484). *Eupatorium album* L., distinguished at a glance by its oblanceolate, narrowly obovate, broadly oblong or narrowly ovate leaves and its prolonged acuminate or attenuate scarious involucre bracts, has proved somewhat perplexing in the field. Upon studying all the material at hand the species seems to resolve itself into four geographic varieties, three of them strongly defined, the fourth perhaps better treated as a forma, though, because of a certain geographic segregation, I am, for the present, maintaining it as a variety. For a beautifully clear photograph of the type of *E. album* I am indebted to the well-known kindness of Mr. Spencer Savage, Assistant Secretary of the Linnean Society of London. As I understand the species it breaks as follows:

- a. Principal leaves spatulate, oblanceolate or narrowly obovate, obtuse, narrowed at base, they and the stem villous or strongly pilose.
 Involucre glandless or essentially so Var. *typicum*.
 Involucre copiously dark-glandular Var. *glandulosum*.
- a. Principal leaves oblong, oblong-lanceolate or narrowly oblong-ovate, acute, the broad bases more rounded, their pubescence short and sparse or wanting.
 Leaves firm, the larger with 10–20 prominent coarse teeth on each margin, pinnately veined Var. *monardifolium*.
 Leaves submembranaceous to firm, the larger with 3–10 low teeth on each margin or entire, triple-nerved from the base Var. *subvenosum*.

Var. **typicum**. *E. album* L. Mant. 111 (1767).—Dry or sandy woods, thickets and clearings, Coastal Plain, New Jersey to Cape Charles, Virginia, locally inland to the District of Columbia; along or on the mountains, South Carolina and Georgia, extending out to the Coastal Plain in South Carolina, northwestern Florida and southeastern Alabama. FIGS. 1 and 2. MAP 35.


In our Virginia field-work we have seen var. TYPICUM (with glandless or essentially glandless involucre) only on the Eastern Shore where it is abundant. We have never met it or any variety of the species in the two southeastern counties; but from Nansemond County to the Fall Line all the plants (very many) which we have seen have copiously glandular involucre and belong to the next.

*Var. **glandulosum** (Michx.) comb. nov. *E. glandulosum* Michx. Fl. Bor.-Am. ii. 98 (1803). *E. stigmatosum* Bertol. Misc. v. 15, t. 5 (1846).—From the Inner Coastal Plain and Piedmont of Maryland and Virginia to central Arkansas, south to northern Florida, southern Alabama, southern Mississippi and (?) Louisiana. FIG. 3. MAP 36.

Michaux's diagnosis, "calycibus 5-floris, creberrime glanduloso-punctatis, lineari-subulatis," supported by a photograph of his type

("in aridis sylvarum Carolinae") secured by the late Dr. Robinson, leaves no doubt as to the identity of his *E. glandulosum*. Similarly Bertolini's beautiful plate and his "squamis . . . linearibus, acuminatis . . . nigropunctatis" satisfactorily identify his *E. stigmatosum*. In our Virginia field-work we did not meet var. *glandulosum* on the Eastern Shore, where var. *typicum* abounds; conversely, although var. *glandulosum* abounds on the Inner Coastal Plain, we met no satisfactory var. *typicum* there.

*Var. **monardifolium**, var. nov. (TAB. 484, FIG. 4-6), foliis oblongis vel oblongo-lanceolatis vel anguste oblongo-ovatis firmis scabris basi rotundatis apice acuminatis margine grosse serratis vel serrato-dentatis, dentibus utrinque 10-20.—Cape May, New Jersey and adjacent Delaware; Prince George County, Maryland to western



MAP 35, range of EUPATORIUM ALBUM var. TYPICUM; 36, var. GLANDULIFERUM; 37, var. MONARDIFOLIUM; 38, var. SUBVENOSUM.

North Carolina. NEW JERSEY: dry, sandy soil, Cape May Point, August 26, 1922, *Fogg*, no. 195. DELAWARE: oak copse, Rehoboth, September 5, 1908, *J. R. Churchill*. MARYLAND: dry soil, between Muirkirk and Contee, September 5, 1910, *A. H. Moore*, no. 4823; gravelly field, near Chillum, September 24, 1926, *S. F. Blake*, no. 9723 (TYPE in Gray Herb.). VIRGINIA: wooded ridge north of Beverley, Fauquier Co., September 29, 1935, *H. A. Allard*, no. 994. NORTH CAROLINA: Burke Co., *Buckley* (cited by Gray in Synop. Fl. as doubtfully his var. *subvenosum*); moist grounds near Biltmore, September 2, 1897, *Biltmore Herb.*, no. 399^b; woodlands near Biltmore, September 14, 1909, *Biltmore Herb.*, no. 3816.^b MAP 37.

When he described *Eupatorium album*, var. *subvenosum* from Long Island and the Pine Barrens of New Jersey Asa Gray cited the Burke County plant with doubt; and on the sheet he noted "Leaves less 3-nerved." This is indeed the case and with the coarser and more abundant teeth and the harsher and firmer leaf-surfaces the pinnate venation well sets off var. *monardifolium*. Its limited and rather inland range, as thus far known, and its isolation at Cape May and in ad-




Photo. E. C. Ogden.

CHRYSOPSIS MARIANA, both figs. $\times 8$: FIG. 1, involucre from New Jersey; FIG. 2, from Virginia.

VAR. *MACRADENIA*, both figs. $\times 8$: FIG. 3, involucre from Waverly, Virginia (TYPE); FIG. 4, from near Williamsburg, Virginia.




Photo. E. C. Ogden.

RUDBECKIA HIRTA: FIG. 4, lower surface of leaf, $\times 10$, from Rhode Island.
 Var. SERICEA: FIG. 3, lower surface of leaf, $\times 10$, from North Dakota.
 Var. CORYMBIFERA: FIG. 1, TYPE, $\times \frac{2}{5}$; FIG. 2, lower surface of leaf, $\times 10$.

jaacent Delaware are interesting in view of the considerable continental element in the Cape May flora, as emphasized by Stone and others.

Var. SUBVENOSUM Gray, Synop. Fl. N. Am. i². 98 (1884).—Known only from pine-barrens of Long Island, New Jersey and Delaware. FIGS. 7 and 8. MAP 38.

*EUPATORIUM LEUCOLEPIS (DC.) Torr. & Gray. CHESTERFIELD Co.: exsiccated argillaceous swale west of Petersburg Turnpike, north of Swift Creek, *F. & L.*, nos. 6408, 6878. PRINCE GEORGE Co.: argillaceous and siliceous boggy depression, about 3 miles southeast of Petersburg, at head of Poo Run, *F. L. & S.*, no. 6879. See pp. 344 and 360.

The first collections, apparently, from between South Carolina and Delaware. See p. 345. The Virginia plant, like the material from South Carolina, Georgia, Florida, Alabama and Louisiana is quite like the typical plant of Delaware and New Jersey. The species was based on *E. glaucescens*, β *leucolepis* DC. Prodr. v. 177 (1836). This was the characteristic New Jersey plant, as shown by a photograph of the type secured by the late Dr. B. L. Robinson in 1905.

The plant of southern Rhode Island and southeastern Massachusetts which has passed as *Eupatorium leucolepis* is a strongly defined geographic variety which I am calling

EUPATORIUM LEUCOLEPIS (DC.) Torr., var. **novae-angliae**, var. nov. (TAB. 485, FIG. 3-5), caulis internodiis superioribus pilosis; foliis plerumque planis subacuminatis acute serratis dentibus subpatis subtus pilosis vel hispidis; venis lateralibus primariis subtus elevatis basi longe decurrentibus.—Pond-shores, Plymouth Co., Massachusetts and Washington County, Rhode Island. MASSACHUSETTS: shore of Smelt Pond, Kingston, August 30, 1908, *W. P. Rich & C. H. Knowlton*; muddy margin, Micajah's Pond, Plymouth, September 13, 1925, *L. B. & F. E. Smith, jr.*; edge of Triangle Pond, Plymouth, August 26, 1928, *Griscom*, no. 12,706 (exceptionally small-leaved); gravelly upper beach of King Pond, Plymouth, August 30, 1928, *Fernald & Griscom*, no. 1076; damp sandy shore of Loon Pond, Lakeville, August 26, 1913, *Fernald & Long*, no. 10,492 (TYPE in Gray Herb.). RHODE ISLAND: sandy and peaty shore, southern end of Long Pond, South Kingstown, September 5, 1914, *Collins & Fernald*, no. 11,444; granitic gravel and sand about small pond east of Long Pond, South Kingstown, September 5, 1914, *Collins & Fernald* in Pl. Exsicc. Grayanae, no. 280.

In typical *Eupatorium leucolepis* (FIGS. 1 and 2) the pubescence of the stem is much finer than in var. *novae-angliae*, a cinereous puberulence (FIG. 2); the leaves are commonly plicate and blunt, with sup-